

The maths and facts behind Australian savanna greenhouse gases

Adapted from Culture, Ecology and Economy of Fire Management in North Australian Savannas REKINDLING THE WURRK TRADITION; by Jeremy Russell-Smith, Peter Whitehead, Peter Cooke; CSIRO Publishing; 2009



How much of Australia is affected by fire?

Many of us living in southern Australia probably believe that we 'southerners' have to endure a large number of bushfires each year. But if we look at what percentage of land is affected by bushfires across Australia we can see that the south experiences a very low number of bushfires compared to the north. The three main regions of Australia that are affected by fire each year are:

- 0.6% of southern Australia
- 5% of central Australia
- 23% of northern Australia

How much of the NT burns?

Between April 2002 and March 2003, fires affected 28.6% of the Northern Territory. These figures include an exceptionally large area of fire in arid central Australia in 2002. In total, 210 000 km² of the Northern Territory's tropical savannas region is affected by fire in an average burning year.

How many people?

The tropical savanna goes across northern Australia and takes up 26% of Australia's land mass. In 2006, 495,000 people lived across the northern Australia. Many of these live in cities and major urban centres. In the rural areas outside the cities there is only one person for every 7km².

How many Indigenous peoples?

18.7% of people in northern Australia are Indigenous Australians. The proportion of the population that is Indigenous is much greater outside the major centres. For example: in the Kimberley, nearly half of the population is Indigenous; in the Northern Territory savannas outside Darwin, it is more than 70%; and in very remote regions it can be more than 90%. All in all, about 19% of the tropical savannas region is presently owned or managed by Indigenous people.

How many Indigenous languages?

The Aboriginal groups (nations) across northern Australia have about 20 language families.

How many cattle?

There are about 5 million cattle being farmed on the tropical savanna.

Page 1.

For more information on how you can help our environment, or to make some suggestions of your own, please go to www.coolaustralia.org



coolaustralia.org

The maths and facts behind Australian savanna greenhouse gases

How much income?

In 2006 an estimate of income for different industries in the tropical savanna was:

- Mining of minerals (excluding coal) - \$6 billion
- Cattle grazing - \$1.4 billion
- Tourism and fishing - \$2 billion

Percentage tropical savanna greenhouse gas emissions?

Approximately 3% of Australia's greenhouse gas emissions result from savanna fires.



Greenhouse gas reductions

In 2008 Professor Ross Garnaut prepared one of the most important climate reports for the Australian government, the Garnaut Climate Change Review. In this report Garnaut estimated that the savanna agricultural emissions from northern Australia could be reduced by 50%.

How have the Earth's greenhouse gases increased?

Gas	1750	Now	Percentage increase
Carbon dioxide	280ppm	390.5ppm	39.5%
Methane	700ppb	1871ppb	16.7%
Nitrous oxide	270ppb	323ppb	19.6%

ppm = parts per million (how many atoms or molecules per million atoms and molecules)

ppb = parts per billion (how many atoms or molecules per billion atoms and molecules)

CO₂-e of greenhouse gases

The CO₂-e is the carbon dioxide equivalent for greenhouse gases. Other greenhouse gases are measured against the warming properties of CO₂ in the atmosphere.

Gas	CO ₂ -e
Carbon dioxide	1
Methane	25
Nitrous oxide	298
CFC-12	10900
HCFC-22	1810
Tetrafluoromethane	11200
Sulfur hexafluoride (SF ₆)	22800

Page 2.

For more information on how you can help our environment, or to make some suggestions of your own, please go to www.coolaustralia.org



coolaustralia.org

The maths and facts behind Australian savanna greenhouse gases

The formula to calculate the amount of nitrous oxide and other nitrous gases is:

And for N₂O and NO_x is:

$$E_{ij} = EF_{ij} A_j^1 P_j \sum_k (FL_{jk} BEF_{jk} CC_{jk} NC_{jk}) M_i$$

Australia's biomass fires

All of Australia's bush (also called biomass) fires contributes about 6–8% of global carbon emissions from biomass fires.

What are the predicted climate changes for northern Australia in 2030?

Predicted climate changes for northern Australia by 2030 due to greenhouse gases are:

- 1 degree increase in temperature.
- More days with temperatures higher than 35 degrees.
- Increase evaporation resulting in the country drying out faster.
- No large changes in the amount of rainfall.
- Greater chance of damaging cyclones and heavy rains.
- Lightning occurs more often.

Damaging cyclones

In 2006, Cyclone Monica, which knocked over more than 50% of trees in 700 km² of savanna forest and woodland in Arnhem Land.

Weeds and burning

Gamba grass is an introduced tall grass from West Africa originally planted as cattle feed. It can reach 4–5 m in height and have a standing biomass of up to about 30 tonne per hectare (about 5–7 times that of the native grasses). When it is that tall and thick, Gamba grass burns much hotter than native grasses.

Impacts on native flora and fauna

Fires in the tropical savanna are already thought to be responsible for the mainland extinction of at least one animal; a small wallaby called a Mala. We can expect further hot burns to have a negative impact on other native plants and animals.



Page 3.

For more information on how you can help our environment, or to make some suggestions of your own, please go to www.coolaustralia.org



coolaustralia.org

The maths and facts behind Australian savanna greenhouse gases

References

Page 317

Table 12.2: A comparison of emission factors from fires in savannas in Australia, Brazil, and southern Africa and forest fire in southern Australia. Northern Australian savanna¹ CO₂ CO CH₄ 0.87 0.078 0.0035 Southern Australian forest² 0.85 0.091 0.0054 Brazilian cerrado³ 0.94 0.05 0.002 1: Hurst et al. 1994a; 2: Hurst et al. 1996; 3: Ward et al. 1992; 4: Ward et al. 1996 Southern African savanna⁴ 0.93 0.065 0.00375

Page 318

Table 12.2: A comparison of emission factors from fires in savannas in Australia, Brazil, and southern Africa and forest fire in southern Australia. Northern Australian savanna¹ CO₂ CO CH₄ 0.87 0.078 0.0035 Southern Australian forest² 0.85 0.091 0.0054 Brazilian cerrado³ 0.94 0.05 0.002 1: Hurst et al. 1994a; 2: Hurst et al. 1996; 3: Ward et al. 1992; 4: Ward et al. 1996 Southern African savanna⁴ 0.93 0.065 0.00375

Pages 331-332

Formula for greenhouse gases

Page 3.

For more information on how you can help our environment, or to make some suggestions of your own, please go to www.coolaustralia.org



coolaustralia.org